

Pre-Knowledge Topics Answers to problems

Q6.1 a) $85.2/284 = 0.3$ moles b) $73.56/122.6 = 0.6$ moles c) $249.5/249.5 = 1.0$ moles

d) $0.125 \times 212.8 = 26.6$ g e) 2Mg : 2O or 1:1 ratio 2.4g of Mg = 0.1moles so we need 0.1 moles of oxygen (O₂): $0.1 \times 32 = 3.2$ g

7.1 a) $9.53\text{g}/95.3 = 0.1$ moles, in 100cm^3 or 0.1dm^3 in 1dm^3 $0.1\text{moles}/0.1\text{dm}^3 = 1.0 \text{ mol dm}^{-3}$

b) $13.284\text{g}/331.2 = 0.04$ moles, in 2dm^3 in 1dm^3 $0.04\text{moles}/2\text{dm}^3 = 0.02 \text{ mol dm}^{-3}$

c) 100cm^3 of $0.1 \text{ mol dm}^{-3} = 0.01$ moles added to a total volume of $2 \text{ dm}^3 = 0.01\text{moles}/2\text{dm}^3 = 0.005 \text{ mol dm}^{-3}$

d) in 1dm^3 of 1 mol dm^{-3} silver nitrate, 1 mole of Ag = 107.9g in $0.1\text{dm}^3 = 107.9 \times 0.1 = 10.79$ g

e) $0.0526 \times 79.7 = 42.0274$ g

=====

8.1

Ba(NO₃)₂ : Na₂SO₄

1 : 1 ratio

12.5cm^3 of Ba(NO₃)₂ = 0.0125dm^3

$0.15 \text{ mol dm}^{-3} \times 0.0125\text{dm}^3 = 0.001875$ moles

same number of moles of sodium sulfate needed, which has a concentration of 0.25 mol dm^{-3}

$0.001875 \text{ moles} / 0.25 \text{ mol dm}^{-3} = 0.0075 \text{ dm}^3$ or 7.5cm^3

=====

9.1 1-chlorobutane



Add butan-1-ol to concentrated HCl and shake

9.2 react ethene with hydrogen gas at high temperature and pressure with a nickel catalyst

The reaction is similar in that it releases hydrogen but different as it proceeds much slower than in water

9.3 propanal

propanone



The carbon atom joined to oxygen in propanal has a hydrogen attached to it, it does not in propanone.


=====

Mark scheme – Topics 6-10

10.1 An acid is a proton donor

10.2 Ammonia can accept a proton, to become NH_4^+

10.3 ethanoic acid has not fully dissociated, it has not released all of its hydrogen ions into the solution.



Mostly this Very few of these

10.4 $\text{pH} = -\log [0.01] = 2$ **The pH = 2**